



1. IF YOUR CHILD HAS HEAD LICE

If your child has head lice it should be treated immediately and checked daily for at least a week later to ensure they are all gone. If lice or eggs are still found, this process will need repeating. If using a chemical treatment always follow the manufacturer's guidance.

If you discover your child has head lice you should keep your child off school to treat them. This will take no longer than two hours, once treated they must return to school.

If you have treated your child for head lice, please inform the school so that there is no confusion, this is because when head lice are found on a child at school, parents will be contacted to collect their child for treatment and then return to school.

If parents/carers keep their child off school for *more than* half a day due to him or her having head lice the absence will not be authorised.

If your child has long hair it is advisable during an outbreak to tie hair back to help prevent infestation.

2. FACTS

- Head lice are small, six-legged wingless insects, pin-head size when they hatch, less than match-head size when fully grown and grey/brown in colour.
- They are difficult to detect in dry hair even when the head is closely inspected.
- They very often cause itching, but this is not always the case, particularly when recently arrived on the head.
- Head lice cannot fly, jump or swim, but spread by clambering from head to head.
- Anyone with hair can catch them, but children who have head to head contact, either at school or during play, are most commonly affected.
- Head lice feed by biting and sucking blood through the scalp of their host.
- The female louse lays eggs in sacs (nits) which are very small, dull in colour, and well camouflaged.
- These are securely glued to hairs where the warmth of the scalp will hatch them out in 7-10 days.
- Empty egg sacs are white and shiny and may be found further along the hair shaft as the hair grows out.
- Lice take 6-14 days to become fully grown, after which they are capable of reproduction.
- Head lice are not fussy about hair length or condition.
- Clean hair is therefore no protection, although regular (e.g.: weekly) hair washing and combing sessions offer a good opportunity to detect head lice, and arrange treatment if discovered.



- There is no need to wash or fumigate clothing or bedding that comes into contact with head lice.

3. DETECTION

Head lice are well camouflaged and hide when disturbed by combing. They do not always cause itching, particularly when recently arrived on the head. They may also be few in number and a quick inspection is unlikely to detect them.

The following method of detection is effective:

Wash the hair using ordinary shampoo, then use a wide toothed comb to straighten and untangle the hair. It is easier to do this with wet hair, and using hair conditioner. Once the comb moves freely through the hair, without dragging, switch to a fine tooth comb. Make sure the teeth of the comb slot into the hair at the roots and draw down to the ends of the hair with every stroke. The comb must be fine enough to catch the lice - size indicated below - do not confuse lice or their eggs with clumps of dandruff or other debris. Check the comb for lice after each stroke as you work through the hair section by section, so that the whole head of hair is combed through, then rinse out the conditioner and repeat the combing procedure in the wet hair. This can be undertaken on a regular basis – e.g.: at routine hair washing sessions - to detect the presence of lice before they can spread. Check all family members at the same time and arrange treatment when lice are found.

If you find lice, then there are two options.

Whichever option you choose it is important to recognise that neither will protect against re-infection if head to head contact is made with someone with head lice at a later date. You may therefore wish to undertake occasional checks during hair washing sessions.

4. TREATMENT

Do not use lotions unless you find a living moving louse. Check all close family/friends by the "wet combing" method, as described, and treat anyone who is found to have lice at the same time, to prevent re-infection. Ensure you have enough lotion to treat all those affected and follow the instructions on the packet carefully, e.g.: as to how long the treatment must remain on the hair to be effective, how often you may apply the product etc. The product may be capable of killing eggs, as well as lice, but there is no certainty of this. Check for baby lice hatching out from eggs 3-5 days after you use it, and again at 10-12 days. If the lice appear to be unaffected by the product (some lice may have developed resistance to a particular insecticide) or if the problem persists then you should take advice from your local school nurse, health visitor, pharmacist or GP, who will be able to advise you on alternative treatments and explain how to use these to best effect.



You should seek advice where whoever is being treated is either under 1 year of age, suffers from asthma or allergies, or is pregnant or breast feeding.

5. FURTHER INFORMATION

Further Information If you are at all worried about head lice or feel you need more advice on how to cope, then you should consult your school nurse, health visitor, pharmacist or family doctor. Information provided by: Department of Health For further information visit: <http://www.dh.gov.uk/health>